

# Knee Arthroscopy: Debridement / Meniscectomy



Knee Arthroscopy: Debridement / Meniscectomy - Physical Therapy Prescription

\*\*\*Please fax progress notes to Advanced Orthopedics & Sports Medicine at 720-895-1121\*\*\*

Name:	Date of Surgery:
Diagnosis: R / L Knee Arthroscopy – meniscectomy, chondroplasty, debridement	
Frequency: 2-3 times per week for weeks,	Therapy to start 3-5 days after surgery

#### POSTOPERATIVE MANAGEMENT

- Reduce pain/swelling (ice 3x's daily for 15-20 minutes directly on knee)
- Full knee ROM as soon as possible
- Full weight bearing as tolerated
- Normal gait (walking) without crutches/cane

### Week 0-2:

- · Modalities as needed
- Stationary bike with high seat if needed and lower it to normal height when able
- Ouad sets/SLR
- Chair Squats/Wall squats keep tibia perpendicular to floor
- Open/Closed chain ex's (leg extensions, leg curls, leg press etc)
- Step Ups start with comfortable height and progress to normal step height as able

## Week 3-4:

- Cont. as above
- Stairmaster, Versa climber, general cardio
- Exercise affected leg only to develop symmetrical LE strength

#### Week 5-6:

- Cont. as above
- May begin running if knee is not swollen or painful
- Plyometrics if needed
- Introduce sport specific training/functional movements

## Criteria for discharge:

- 1. Full, pain free range of motion
- 2. Strength is equal bilaterally

Stephen G. Thon, MD

- 3. Has met specific functional/activity goals
- 4. Has been cleared by physician

Signature:	www.AdvancedOrtho.org
Date:	Phone: (303) 344-9090

Fax: (720) 895-1121